

11TH INTERNATIONAL CONFERENCE
ON
SIMULATION AND MODELLING
IN THE
FOOD AND BIO-INDUSTRY
2020

FOODSIM'2020

EDITED BY

Jan F.M. Van Impe

and

Monika E. Polańska

SEPTEMBER 6-10, 2020

GHENT, BELGIUM

A Publication of EUROSIS-ETI

Printed in Ghent, Belgium

Cover picture Cain and Abel - detail of The Adoration of the Mystic Lamb (brothers Hubert and Jan van Eyck) is licensed under the Creative Commons Attribution-Share Alike.

MICROBIAL DETERIORATION OF LAMB MEAT OF PORTUGUESE ORIGIN AS AFFECTED BY ITS INTRINSIC PROPERTIES

Vasco A. P. Cadavez
Ursula Gonzales-Barron
Diogo Félix-Oliveira
Sara Coelho-Fernandes
Gisela Santos-Rodrigues

Centro de Investigação de Montanha (CIMO)
Instituto Politécnico de Bragança
5300-253 Bragança, Portugal
vcadavez@ipb.pt

José M. Lorenzo
Roberto Bermúdez Piedra

Fundación Centro Tecnológico da Carne
Parque Tecnológico de Galicia
3200 San Cibrao das Viñas, Spain
jmlorenzo@ceteca.net

KEYWORDS

Sheep, local breeds, proximate composition, pH, lactic acid bacteria, *Pseudomonas*, psychrotrophic bacteria, shelf-life

ABSTRACT

In Portugal, sheep and goat meat production constitutes 2.8% of the total meat production, with a self-sufficiency of ~82%. The main autochthonous sheep breeds exploited for meat production are Churra-Galega-Bragançana (CGB) and Bordaleira-de-Entre-Douro-e-Minho (BEDM), whose quality must be optimised in order to ensure adequate income levels for sheep producers. The study aimed to characterise the evolution of spoilage microorganisms in refrigerated vacuum-packed (VP) lamb meat from BEDM and CGB breeds; and elucidate how intrinsic properties of meat can affect its microbial spoilage. Meat from BEDM breed presented higher ($p < .0001$) populations of mesophiles, lactic acid bacteria, *Pseudomonas* spp. and psychrotrophic bacteria, since its higher ultimate pH (means: 5.77 for BEDM vs. 5.58 for CGB) accelerated spoilage rate ($p < .0001$). While water activity and protein content were not found to modulate microbial deterioration ($p > 0.05$), the growth of spoilage bacteria was found to be exacerbated by higher moisture ($p < .0001$) and higher ash content ($p < 0.001$). By contrast, a higher fat content retarded ($p < .0001$) the growth of spoilage bacteria in VP lamb meat. In order to extend the shelf-life of Portuguese-origin lamb meat, animal handling must be enhanced to minimise pre-slaughter stress, and a carcass classification system should be adopted towards the selection of fatter animals and chilled carcasses of optimal ultimate pH.

INTRODUCTION

Sheep farming is an activity of high economic importance as these animals are producers of meat, milk, wool and fur. In Portugal, sheep and goat meat production constitutes 2.8% of the country's meat production (INE 2016), with a level of self-sufficiency of 82% (Santos-Rodrigues et al. 2019). Therefore, increasing the production of sheep meat and optimising its quality, making it more attractive to consumers, is essential to ensure a good income level for sheep producers.

Portuguese sheep production is featured by a large diversity of production systems, all based on the use of natural pastures and agricultural residues (Cruz et al. 2019). These

animal production grazing systems are characterised by great heterogeneity and have contributed to the maintenance of the ecological balance of the natural pastures, and the environmental and socio-cultural richness of the landscapes of Portugal. The sixteen national indigenous sheep breeds, in addition to contributing to the diversity of production systems, are an important genetic heritage that must be preserved. Some of these breeds, classified as threatened, have small body size and good adaptation to adverse environments (climate and orography), which makes them particularly well suited to the use and enhancement of natural pastures. Two of the autochthonous sheep breeds exploited for meat production are Churra-Galega-Bragançana (CGB) and Bordaleira-de-Entre-Douro-e-Minho (BEDM), from the Mediterranean bioregion and the Atlantic bioregion, respectively (Cruz et al. 2019).

Today's consumer is sensitive to management practices capable of improving animal welfare, and is willing to pay a higher price for quality certified meat. Enhancing the quality of meat of autochthonous breeds could contribute to the preservation of the rural world and its diversity, as well as increasing the profitability of these production systems. In this way, we will be able to ensure the conservation of endangered breeds and improve the living standards of the sheep farmers that remain in the rural areas of Portugal.

However, meat quality is a multifactorial concept regulated by factors that are intrinsic and extrinsic to the animal. On the one hand, quality is perceived by the consumer through good sensorial properties – optimised through cold maturation; while, on other hand, the producer must not only meet such organoleptic demands, but must also ensure that the product remains safe during its life time, which should aim to be the longest possible to avoid economic losses (Mills et al. 2014).

In the case of lamb meat, after 7 days of maturation ~80% of its maximum tenderness potential is reached (Prates 2000). However, during this maturation, microbial deterioration takes place due to the proliferation of psychrotrophic bacteria, lactic acid bacteria, *Pseudomonas* spp., *Clostridium* spp., etc. (Clemens et al. 2010). One of the well-known strategies to prolong the life of the meat is vacuum packaging, which can moderately retard microbial deterioration. However, the extent of such retardation depends on chilling system/profile, initial microbial contamination and the physicochemical or intrinsic properties of meat.

Therefore, the objective of this study was twofold: (i) to evaluate the evolution of spoilage indicator microorganisms (mesophiles, psychrotrophic, lactic acid bacteria and *Pseudomonas* spp) in refrigerated vacuum-packed lamb meat from two Portuguese breeds, BEDM and CGB; and (ii) to elucidate, by means of mixed models, any interrelationship between meat's intrinsic properties (i.e., pH, water activity [a_w], moisture content, fat content, protein content and ashes) and microbial growth.

METHODOLOGY

Lamb rearing and feeding

In the Mediterranean region, located in Bragança, CGB lambs were raised on the holding of the School of Agriculture of the Polytechnic Institute of Bragança. In the Atlantic bioregion, located in Ponte de Lima, BEDM lambs were raised on the holding of the Ponte de Lima Agrarian School. In both bioregions, the production system used was the semi-intensive one, whose feeding was based on grazing on natural pastures. The hours of grazing varied according to hours of light, heat and herd size. In winter, the flocks would be released in the morning to graze all day until dark. In the summer, the herds would leave at dawn and graze until midmorning, then they would be put in a stable under shade, and would come out when the heat had subsided. Once on the premises, lambs had access to meadow hay, supplemented with protein and mineral-rich concentrates. In the stable, the lambs had water *ad libitum*. The lambs were not weaned, and were reared in the fall of 2018 and spring 2019. For this investigation, 15 BEDM and 15 CGB lambs were reared in 2018 and 15 BEDM and 15 CGB lambs were reared in 2019.

Preparation of meat samples

The slaughter and therefore the analyses were carried out in batches in the same abattoir: in 2018, three batches of 5 CGB lambs each, and three batches of 5 BEDM lambs of 5 each were slaughtered; and in 2019 two batches of 7 and 8 CGB lambs each, and two batches of 7 and 8 BEDM lambs each were slaughtered.

All lambs were four months old when slaughtered, and the carcasses obtained were chilled at 4 °C for 24 hours. After carcass splitting, the *Longissimus dorsi* muscle was removed from the 6th to the 13th vertebra under aseptic conditions. The left side was divided into three parts. Each of them was vacuum packed (Silvercrest SFS 110 B2, Germany); labelled with the number 3, 9 or 15, corresponding to the day of microbiological analysis; and stored at 4±0.5 ° C (Portiso ECB-3000, UK). Concentrations of total viable counts, psychrotrophic bacteria, lactic acid bacteria and *Pseudomonas* spp. were determined at each time point.

The right half of the *L. dorsi* muscle was kept for the physicochemical analyses – pH, moisture and dry matter, fat content, protein content and ash content – which were carried out on day 1 after slaughter. Only a_w was measured on day 3 post-slaughter. Unlike microbiological analyses, the *intrinsic properties of meat* (physicochemical analyses) were assayed right at the beginning of the cold maturation.

Microbiological and physicochemical analyses

For the microbiological analyses, twenty-five g of meat were homogenised in 225 ml of buffered peptone water (611014 Liofilchem, Roseto degli Abruzzi, Italy) for 1 min (Interscience Bag Mixer 400, France). One-ml aliquots from decimal dilutions were inoculated on Aerobic Count Plate petrifilms (3M, MN, USA) for counting of mesophiles, and on Lactic Acid Count Plate petrifilms (3M, MN, USA) for counting of lactic acid bacteria. One-ml aliquots were plated by incorporation in Plate Count Agar (610040 Liofilchem, Roseto degli Abruzzi, Italy) for the determination of psychrotrophic bacteria, while, for the quantification of *Pseudomonas*, 0.5-ml aliquots were spread onto *Pseudomonas* Agar Base (CM0559 Oxoid, Thermo Fisher Scientific, UK), added with 1% v/v glycerol and supplemented with cefrimide-fucidine-cephalosporin (610071 Liofilchem, Roseto degli Abruzzi, Italy). The mesophilic and lactic acid bacteria plates were incubated at 35±0.5°C for 48 h; the psychrotrophic bacteria plates at 7±0.5°C for 11 days; and the *Pseudomonas* plates at 25±0.5°C for 24 h. Plating was done in duplicate, and colony counts were transformed to log CFU/g.

The meat's intrinsic properties measured were pH, a_w and proximate composition. The pH measurement was carried out according to Franco (2009), using a pH meter (HI 99163, Hanna Instruments, Eibar, Spain) equipped with a 232D glass penetration probe. To measure a_w , lamb steaks were cut to exactly fit in the cuvette of the Aqualab meter (4TE Decagon, USA). a_w was recorded after measurement stabilisation. Moisture, fat, protein and ashes contents were determined according to ISO (1997), AOCS (2005), ISO (1978) and ISO (1998), respectively. Determinations were made in triplicate per meat sample. Contents of fat, protein and ashes were expressed in dry basis.

Mixed-effects models

The statistical analysis aimed to understand to what extent the intrinsic properties of meat (i.e., animal breed, pH, a_w and proximal composition) can affect or modulate its microbial spoilage, as characterised by the change in the populations of mesophiles, lactic acid bacteria, psychrotrophic bacteria and *Pseudomonas* spp.

A general mixed-effects model of the type,

$$Y_{rj} = \beta_{0j} + \beta_{1r} + \beta_2 Day + \beta_3 (X)(Day) + \beta_4 Day^2 + \varepsilon_{rj}$$

$$\beta_{0j} = \beta_0 + v_j \quad \dots(1)$$

was adjusted to each of the microbial groups (Y) to assess the effect of breed, time of maturation and each of the intrinsic properties of meat (X), in separate. The response variable Y_{rj} is the microbial concentration in the meat sample from lamb j of breed r , measured after a maturation time Day . The model intercept β_0 is affected by random shifts v_j caused by the different lambs j . These random effects were removed from the model assuming they follow a normal distribution. The parameter β_{1r} is the fixed effect of lamb breed, while β_2 the linear effect of maturation time.

It is allowed that the effect of maturation time could be modulated by the intrinsic property X , so the parameter β_3 assesses this interaction. A quadratic effect for Day (β_4) was added given its significance in all models. The residuals ε_{rj} are assumed to follow a normal distribution.

Equation (1) was adjusted separately to each of the four microbial groups (Y), and in each of the adjustments, the independent variable X represents pH, a_w , humidity, protein, fat or ashes content. Thus, in total 24 models were fitted. The results of the mixed linear models presented in the following Section include: (i) parameter estimates along with standard errors; and (ii) the significance of the sources of variation tested in analysis of variance (ANOVA) by the F-test. The models were adjusted in the R software (The R Core Team 2019).

RESULTS AND DISCUSSION

If lamb is produced under good manufacturing practices, the initial counts of microorganisms on the meat surface are likely to be $\sim 10^3$ CFU/cm² or lower (Mills et al. 2014). In our experiments, the abattoir's controlled process hygiene ensured that the bacterial counts in vacuum-packed meat were relatively low still on the third day after slaughter, at mean values of 1.45 log CFU/g for mesophiles, 0.899 log CFU/g for LAB, 1.27 log CFU/g for psychrotrophic bacteria and 1.02 log CFU/g for *Pseudomonas* spp (Figure 1). By contrast, the mesophiles level determined by Wang et al. (2019) in VP lamb meat on the third day of storage was much higher at 4.95 log CFU/g despite the comparable storage temperature of 4-6°C. In our work, the lowest initial microbial populations were found for *Pseudomonas* and LAB. Since vacuum packaging excludes oxygen, the strictly aerobic rapidly-growing *Pseudomonas* are inhibited. Mills et al. (2014) explained that, after vacuum-packaging, the population of LAB is generally low (10 CFU/cm²) but it increases during storage until growth stops due to substrate depletion. At -1.5°C, growing LAB populations have been shown to be displaced by succeeding populations without a decline in observable LAB numbers (Jones 2004).

The difference in bacterial population size between lamb breeds was a recurrent fact in the analysed slaughter batches from both years, 2018 and 2019. While deteriorating bacteria increased as maturation took place ($p < .0001$ for the terms Day and Day² in Tables 1-4), the microbial growth trends were clearly different between lamb breeds (Figure 1), which was corroborated by the significant effect of Breed on mesophiles counts ($p < .0001$; Table 1), LAB counts ($p < .0001$; Table 2), *Pseudomonas* counts (p between 0.004 and 0.006; Table 3) and psychrotrophic bacteria counts (p between 0.009 and 0.017; Table 4). In addition, for the four bacterial groups, the increase in population appeared less pronounced in VP meat from CGB lambs (Figure 1), which was also suggested by the negative sign of the model term "Breed - CGB", displayed in Tables 1-4. The fact that the microbial populations of CGB lamb meat were for all microbial groups significantly lower than those of BEDM lamb meat are linked to the lower ultimate pH of the CGB lamb meat (5.58 for CGB versus 5.77 for BEDM; Figure 2).




Figure 1: Increase in Microbial Populations in Vacuum-Packed Lamb Meat Stored at 4°C During 15 Days, by Breed Churra-Galega-Bragança (CGB) and Bordaleira-entre-Douro-e-Minho (BEDM)




Figure 2: Box Plots of Lamb *Longissimus dorsi* pH Measured at 24 hours Post-Slaughter by Breed, Churra-Galega-Bragançana (CGB) and Bordaleira-entre-Douro-e-Minho (BEDM)

Table 1: Effects of Lamb Breed and Initial Intrinsic Factors of Meat on the Concentration of **Mesophilic Bacteria** in Refrigerated Vacuum-Packed Meat as Quantified by Six Separate Linear Mixed Models (F-Values and Associated p-Values from ANOVA are Shown; Quadratic Effect of Day is Significant in All Models but Not Shown)

Model	Term	Estimate (SE)	F (p-value)
pH	Intercept	1.576 (0.337)	533.8 (<.0001)
	Breed - CGB	-0.262 (0.192)	7.503 (0.008)
	Day	-3.598 (1.249)	168.4 (<.0001)
	pH×Day	0.592 (0.209)	7.951 (0.006)
a_w	Intercept	1.689 (0.343)	516.5 (<.0001)
	Breed - CGB	-0.480 (0.178)	7.259 (0.009)
	Day	2.928 (14.04)	159.9 (<.0001)
	a_w ×Day	-3.175 (14.14)	0.050 (0.823)
Moisture (%)	Intercept	1.543 (0.329)	723.0 (<.0001)
	Breed - CGB	-0.196 (0.160)	10.16 (0.002)
	Day	-9.793 (1.813)	169.5 (<.0001)
	Moisture×Day	0.125 (0.023)	28.97 (<.0001)
Fat (% db)	Intercept	1.469 (0.319)	662.8 (<.0001)
	Breed - CGB	-0.055 (0.174)	9.314 (0.004)
	Day	0.133 (0.345)	184.67 (<.0001)
	Fat×Day	-0.066 (0.012)	33.30 (<.0001)
Protein (% db)	Intercept	1.655 (0.342)	515.4 (<.0001)
	Breed - CGB	-0.414 (0.187)	7.243 (0.009)
	Day	-1.804 (1.388)	161.9 (<.0001)
	Protein×Day	0.018 (0.015)	1.392 (0.241)
Ashes (% db)	Intercept	1.578 (0.328)	643.8 (<.0001)
	Breed - CGB	-0.264 (0.166)	9.047 (0.004)
	Day	-1.427 (0.433)	172.0 (<.0001)
	Ashes×Day	0.222 (0.047)	22.25 (<.0001)

The effect of pH on the steeper or slower microbial growth can be also deduced from the interaction term pH×Day, which was significant for the four bacterial groups, namely mesophiles (p=0.006 in Table 1), LAB (p=0.001 in Table 2), *Pseudomonas* (p=0.050 in Table 3) and psychrotrophic

bacteria (p=0.010 in Table 4). The positive estimates for the interaction pH×Day imply that a higher ultimate pH of meat tends to accelerate the microbial growth. Thus, the rate of bacterial population growth is regulated by the ultimate pH.

Table 2: Effects of Lamb Breed and Initial Intrinsic Factors of Meat on the Concentration of **Lactic Acid Bacteria** in Refrigerated Vacuum-Packed Meat as Quantified by Six Separate Linear Mixed Models (F-Values and Associated p-Values from ANOVA are Shown; Quadratic Effect of Day is Significant in All Models but Not Shown)

Model	Term	Estimate (SE)	F (p-value)
pH	Intercept	1.040 (0.311)	300.5 (<.0001)
	Breed - CGB	-0.355 (0.155)	17.70 (<.0001)
	Day	-3.692 (1.085)	56.99 (<.0001)
	pH×Day	0.633 (0.018)	12.24 (0.001)
a_w	Intercept	1.162 (0.325)	311.1 (<.0001)
	Breed - CGB	-0.588 (0.137)	18.32 (<.0001)
	Day	5.347 (11.80)	50.79 (<.0001)
	a_w ×Day	-5.466 (11.88)	0.211 (0.646)
Moisture (%)	Intercept	1.075 (0.316)	340.0 (<.0001)
	Breed - CGB	-0.421 (0.140)	20.02 (<.0001)
	Day	-5.735 (1.653)	53.86 (<.0001)
	Moisture×Day	0.074 (0.021)	12.23 (<.0001)
Fat (% db)	Intercept	1.016 (0.309)	335.4 (<.0001)
	Breed - CGB	-0.308 (0.148)	19.75 (<.0001)
	Day	0.157 (0.340)	56.73 (<.0001)
	Fat×Day	-0.044 (0.010)	17.84 (<.0001)
Protein (% db)	Intercept	1.115 (0.322)	311.0 (<.0001)
	Breed - CGB	-0.499 (0.145)	18.32 (<.0001)
	Day	-2.233 (1.186)	52.31 (<.0001)
	Protein×Day	0.025 (0.013)	3.612 (0.060)
Ashes (% db)	Intercept	1.081 (0.309)	318.3 (<.0001)
	Breed - CGB	-0.432 (0.142)	18.74 (<.0001)
	Day	-0.949 (0.405)	56.67 (<.0001)
	Ashes×Day	0.161 (0.042)	14.67 (0.001)

Meat pH has been shown to affect the growth of bacteria in two ways (Gill 2004). Firstly, the growth of some bacteria is reduced, or inhibited completely, when the pH falls below a certain level. Secondly, the glycolytic processes that determine the ultimate pH also determine the concentration of residual glucose in the meat, and therefore the point at which this preferred growth substrate becomes exhausted and aminoacids start to be metabolised by LAB and *Enterobacteriaceae*, thereby resulting in spoilage. For instance, it is generally accepted that *Enterobacteriaceae* and *Bacillus thermosphacta* are inhibited by pH values lower than 5.8 (Bell 2001).

Although, generally, lamb meat has a higher mean ultimate pH (5.6 – 5.7) than beef meat (pH = 5.5), still the significant effect of Breed in all models (Tables 1-4) suggest that BEDM lambs may have been more susceptible to pre-mortem stress than CGB lambs. The rapid depletion of glycogen levels, prompted by stress, prevents the normal

drop in pH to optimal levels. As a result, meat of higher pH (>5.7) has better conditions for microbial growth – as already explained – ending up ultimately (and unavoidably) in a reduction of shelf-life in refrigerated conditions even when vacuum-packaging is applied.

Table 3: Effects of Lamb Breed and Initial Intrinsic Factors of Meat on the Concentration of *Pseudomonas pp.* in Refrigerated Vacuum-Packed Meat as Quantified by Six Separate Linear Mixed Models (F-Values and Associated p-Values from ANOVA are Shown; Quadratic Effect of Day is Significant in All Models but Not Shown)

Model	Term	Estimate (SE)	F (p-value)
pH	Intercept	1.255 (0.510)	206.3 (<.0001)
	Breed - CGB	-0.369 (0.222)	8.147 (0.006)
	Day	-3.239 (1.657)	32.15 (<.0001)
	pH×Day	0.533 (0.270)	3.795 (0.050)
a _w	Intercept	1.356 (0.513)	215.7 (<.0001)
	Breed - CGB	-0.567 (0.194)	8.520 (0.005)
	Day	-30.48 (17.26)	31.28 (<.0001)
	a _w ×Day	30.50 (17.38)	3.082 (0.082)
Moisture (%)	Intercept	1.188 (0.485)	224.2 (<.0001)
	Breed - CGB	-0.242 (0.203)	8.854 (0.004)
	Day	-11.15 (2.448)	35.53 (<.0001)
	Moisture×Day	0.143 (0.031)	21.00 (<.0001)
Fat (% db)	Intercept	1.115 (0.483)	221.7 (<.0001)
	Breed - CGB	-0.100 (0.215)	8.754 (0.005)
	Day	0.191 (0.534)	36.00 (<.0001)
	Fat×Day	-0.073 (0.015)	22.01 (<.0001)
Protein (% db)	Intercept	1.303 (0.513)	207.5 (<.0001)
	Breed - CGB	-0.464 (0.210)	8.196 (0.006)
	Day	-2.646 (1.785)	31.59 (<.0001)
	Protein×Day	0.028 (0.019)	2.085 (0.151)
Ashes (% db)	Intercept	1.255 (0.501)	217.5 (<.0001)
	Breed - CGB	-0.371 (0.203)	8.591 (0.005)
	Day	-1.288 (0.646)	33.11 (<.0001)
	Ashes×Day	0.201 (0.063)	10.32 (0.002)

In relation to meat a_w, this intrinsic property was not found to modulate the growth of any microbial group, as deduced by the non-significance of the interaction term a_w×Day (p=0.823 for mesophiles; p=0.646 for LAB; p=0.082 for *Pseudomonas* and p=0.315 for psychrotrophic bacteria in Tables 1 to 4, respectively). The lack of effect of a_w is not surprising since vacuum packaging prevents drying at the meat surface, and moisture from within the meat allows the surface a_w to equilibrate to above 0.98. Consequently, there is no inhibitory effect on bacteria once the meat has been packed and stored (Bell 2001). The mean a_w of VP meat measured on the third day after slaughter was the same, 0.9927, for both CGB and BEDM breeds.

Unlike a_w, the growth of spoilage bacteria was found to be exacerbated by the moisture content of meat, as implied by the interaction Moisture×Day that was significant in all bacterial groups (p<.0001 in Tables 1-4). The positive sign

of this interaction suggests that a higher moisture content in VP lamb meat prompted a faster increase in spoilage bacterial numbers. Meat samples may have had different levels of moisture, since lamb carcasses were held for 24 hours in a chilling room at 90% RH with loadings that varied from batch to batch. Under these conditions, moisture loss from lamb carcasses have been reported to be up to 2.2% (Brown et al. 1993). The mean moisture content of lamb meat originating from the BEDM breed was 77.03% (SD=1.517%) whereas that of lamb meat originating from the CGB breed was 75.89% (SD=0.962%).

Table 4: Effects of Lamb Breed and Initial Intrinsic Factors of Meat on the Concentration of *Psychrotrophic Bacteria* in Refrigerated Vacuum-Packed Meat as Quantified by Six Separate Linear Mixed Models (F-Values and Associated p-Values from ANOVA are Shown; Quadratic Effect of Day is Significant in All Models but Not Shown)

Model	Term	Estimate (SE)	F (p-value)
pH	Intercept	1.463 (0.455)	319.6 (<.0001)
	Breed - CGB	-0.251 (0.251)	6.100 (0.017)
	Day	-4.615 (1.686)	159.1 (<.0001)
	pH×Day	0.741 (0.282)	6.893 (0.010)
a _w	Intercept	1.598 (0.464)	330.9 (<.0001)
	Breed - CGB	-0.531 (0.223)	6.308 (0.015)
	Day	18.03 (18.23)	148.7 (<.0001)
	a _w ×Day	-18.54 (18.36)	1.019 (0.315)
Moisture (%)	Intercept	1.387 (0.427)	374.4 (<.0001)
	Breed - CGB	-0.141 (0.221)	7.141 (0.009)
	Day	-13.70 (2.447)	178.5 (<.0001)
	Moisture×Day	0.175 (0.031)	30.76 (<.0001)
Fat (% db)	Intercept	1.288 (0.422)	365.3 (<.0001)
	Breed - CGB	0.045 (0.233)	6.973 (0.011)
	Day	0.134 (0.456)	185.5 (<.0001)
	Fat×Day	-0.091 (0.016)	34.51 (<.0001)
Protein (% db)	Intercept	1.532 (0.462)	324.7 (<.0001)
	Breed - CGB	-0.413 (0.237)	6.192 (0.016)
	Day	-3.261 (1.866)	152.2 (<.0001)
	Protein×Day	0.033 (0.021)	2.576 (0.111)
Ashes (% db)	Intercept	1.468 (0.442)	347.5 (<.0001)
	Breed - CGB	-0.274 (0.226)	6.629 (0.013)
	Day	-1.800 (0.587)	166.1 (<.0001)
	Ashes×Day	0.263 (0.064)	16.87 (0.001)

Interestingly, lamb meat samples of higher fat content underwent a slower microbial deterioration, as implied by the negative interaction Fat×Day that was significant for all microbial groups (p<.0001; Tables 1-4). On the contrary, lamb meat samples of higher content of metal salts and trace minerals (ashes) tended to have a faster microbial spoilage (p of at least 0.001; Tables 1-4). Despite protein content of lamb meat presented a wide range of variation, between 80.97 to 91.34% (db), protein content could not be found to regulate the growth of spoilage bacteria in VP lamb meat. Notice that the interaction term Protein×Day was non-significant (p>0.05) in all bacterial groups (Table 1-4).

CONCLUSION

In order for native-breed lamb meat producers to optimise and meet the challenges posed by the heterogeneity of production chain actors, as well as the challenges of homogeneous quality meat sought by consumers, it is necessary to understand how the intrinsic properties of carcass and meat regulate meat microbiological attributes. This study showed that populations of spoilage bacterial groups were higher in vacuum-packed lamb meat originating from BEDM breed than in that of CGB breed, since ultimate pH was significantly higher in BEDM lamb meat. In addition, a high ultimate pH was demonstrated to increase the rate of microbial deterioration. Other meat intrinsic properties that increased the rate of microbial spoilage were high moisture and ash content. By contrary, a higher total fat content in meat tended to retard its microbial spoilage. In order to extend the shelf-life of Portuguese-origin lamb meat, animal handling and carcass classification can be improved towards the selection of fatter animals and chilled carcasses of optimal ultimate pH.

ACKNOWLEDGEMENTS

The authors are grateful to the Foundation for Science and Technology (FCT, Portugal) and FEDER under Programme PT2020 for financial support to CIMO (UIDB/00690/2020). Dr. Gonzales-Barron acknowledges the national funding by FCT, P.I., through the Institutional Scientific Employment Programme contract.

REFERENCES

- AOCS. 2005. *Official Procedure, Approved Procedure. Am 5-04, Rapid Determination of Oil/Fat Utilizing High Temperature Solvent Extraction*. American Oil Chemistry Society, Urbana, IL.
- Bell, R. G. 2001. "Meat packaging: protection, preservation and presentation". In *Meat Science and Applications*, Y.H. Hui, W.K. Nip, R.W. Rogers & O.A. Young (Eds). Marcel-Dekker, USA, 463-490.
- Brown, T., K. N. Chourouzidis and A.J. Gigiel. 1993. "Spray chilling of lamb carcasses". *Meat Science* 34, 301-304.
- Clemens, R.M., K.H. Adam and G. Brightwell. 2010. "Contamination levels of *Clostridium estertheticum* spores that result in gaseous spoilage of vacuum-packaged chilled beef and lamb meat". *Letters in Applied Microbiology* 50, 591-596.
- Cruz, B. C., J. Cerqueira, J. Araújo, U. Gonzales-Barron and V. Cadavez. 2019. "Estudio de las características de crecimiento de corderos de las razas Churra Galega-Bragançana y Bordaleira-de-Entre-Douro-e-Minho". In *Resúmenes de XVIII Jornadas sobre Producción Animal* (Zaragoza, Spain, May 7-8). Asociación Interprofesional para el Desarrollo Agrario, Spain, 66-68.
- Franco, D., E. Bispo, L. Gonzalez, J.A. Vazquez and T. Moreno. 2009. "Effect of finishing and ageing time on quality attributes of loin from the meat of Holstein-Friesian cull cows." *Meat Science* 83, 484-491.
- Gill, C.O. 2004. "Spoilage, factors affecting – Microbiological". In *Encyclopedia of Meat Science*, W. Jensen (Ed). Elsevier, Oxford, 1324-1330.
- INE. 2016. "Estatísticas Agrícolas 2015". Technical Report, INE, I. P., Instituto Nacional de Estatística, Lisbon, Portugal.
- ISO. 1978. "Determination of Nitrogen Content. ISO 937:1978 Standard". In *International Standards Meat and Meat Products*. International Organization for Standardization. Geneva, Switzerland.
- ISO. 1997. "Determination of Moisture Content. ISO 1442:1997 Standard". In *International Standards Meat and Meat Products*, International Organization for Standardization, Geneva, Switzerland.
- ISO. 1998. "Determination of Ash Content. ISO 936:1998 Standard". In *International Standards Meat and Meat Products*. International Organization for Standardization. Geneva, Switzerland.
- Jones, R. 2004. "Observation on the succession dynamics of lactic acid bacteria populations in chill-stored vacuum-packaged beef". *International Journal of Food Microbiology* 90, 273-282.
- Mills, J., A. Donnison and G. Brightwell. 2014. "Factors affecting microbial spoilage and shelf-life of chilled vacuum-packed lamb transported to distant markets: a review". *Meat Science* 98, 71-80.
- Santos-Rodrigues, G., J.M. Lorenzo, U. Gonzales-Barron and V. Cadavez. 2019. "Características físico-químicas y ternura de la carne de cordero de dos razas autóctonas portuguesas". In *Resúmenes de XVIII Jornadas sobre Producción Animal* (Zaragoza, Spain, May 7-8). Asociación Interprofesional para el Desarrollo Agrario, Spain, 525-527.
- Prates, J. A. M. 2000. "Maturação da carne dos mamíferos: Caracterização geral e modificações físicas". *Revista Portuguesa de Ciências Veterinárias* 95, No. 533, 34-41.
- Wang, T., H. Guo, H. Zhang, F. Ren, M. Zhang, S. Ge, H. Luo and L. Zhao. 2019. "Dynamics of bacterial communities of lamb meat packaged in air and vacuum pouch during chilled storage". *Food Science and Animal Resources* 39, No. 2 (Apr), 209-221.

AUTHOR BIOGRAPHIES

VASCO CADAVEZ, obtained his PhD in Animal Science Engineering at the Portuguese University of Trás-os-Montes and Alto Douro. He is Coordinator Professor in the Animal Science Department of the Polytechnic Institute of Bragança, Portugal (IPB); and has wide expertise in general linear models, linear and non-linear mixed models, count data models, dynamic models and optimisation, meta-analysis and chemometrics statistics which he has applied in the fields of animal science, agriculture and food safety. Dr. Cadavez has published 60 peer-reviewed papers; has disseminated his research at >50 conferences; has organised and taught at 12 international workshops/symposia; and has supervised 20 postgraduate students. He is Editor-in-Chief of the Portuguese Journal *Revista Portuguesa de Zootecnia*.

URSULA GONZALES-BARRON, a Peruvian-Irish national, is a Food Engineer (Honours graduated) with a PhD in Biosystems Engineering from University College Dublin (UCD), Ireland. After a postdoctoral fellowship at the University of Manchester (UK), she became Senior Researcher at UCD, and is currently a Principal Investigator at the CIMO Mountain Research Centre, IPB, Portugal. She leads the "Food Quality and Safety Analytics" research team. Her current research focuses on predictive microbiology, meta-analysis, risk assessment and Bayesian statistics. She is also a Visiting Professor at the National Agricultural University La Molina, Peru. She has published 80 peer-reviewed articles, and has an H-index >20. She acts as Editor of the journal LWT, and speaks fluently English, Portuguese and Spanish.